Fiji

Population	848 000
GNI/capita	4 370 US\$
% urbanization	51%
% people living in cities greater than 100 000 inhabitants	25%
Population below the poverty line (national)	NA
Population below the poverty line (international, <\$1/day)	NA

Under age 5 mortality rate 18/1000 live births (2006) Life expectancy 69 years (2006)

Environmental burden of disease for selected risk factors, per year	r
Estimates based on national exposure and WHO country health statistics 2004	

Risk factor	Exposure		Deaths /year	DALYs/ 1000 cap /year
Water, sanitation and hygiene	Improved water:	47%	<100	1.5
(diarrhoea only)	Improved sanitation:	72%		
Indoor air	SFU% households:	40%	<100	1.2
Outdoor air	Mean urban PM10: 17	ug/m3	-	-
Main malaria vectors	No transmission			
Main other vectors	Aedes fijiensis, Aedes pseudoscutellaris, Aed			Aedes

Environmental burden of disease (preliminary), per year

Estimates based on Comparative Risk Assessment, evidence synthesis and expert evaluation for regional exposure and WHO country health statistics 2004

DALYs/1000 cap	(World - lowest: 13, highest: 289)	31
Deaths		1 100
% of total burden		18%

Environmental burden by diseas		/s/1000 capita], per	year	
	World's		World's	
Disease group	lowest	Country rate	highest	
	country rate		co	untry rate
Diarrhoea	0.2		1.6	107
Respiratory infections	0.1		2.4	71
Malaria	0.0		-	34
Other vector-borne diseases	0.0		0.9	4.9
Lung cancer	0.0		0.0	2.6
Other cancers	0.3		1.4	4.1
Neuropsychiatric disorders	1.4		1.9	3.0
Cardiovascular disease	1.4		5.6	14
COPD	0.0		0.6	4.6
Asthma	0.3		1.2	2.8
Musculoskeletal diseases	0.5		1.4	1.5
Road traffic injuries	0.3		0.7	15
Other unintentional injuries	0.6		4.4	30
Intentional injuries	0.0		0.3	7.5

Other indicators		
Use of leaded gasoline	No (2008)	
Overcrowding	NA	
Malnutrition (% stunting)	NA	

